

Long Wavelength Anomalous Diffusion Mode in the 2D XY Dipole
Magnet.

Ar.Abanov¹, A.Kashuba^{1,2,3} and V.L.Pokrovsky^{1,3}.

Department of Physics, Texas A&M University

College Station, Texas 77843-4242, USA¹

and

Laboratorium für Festkörper Physik der ETHZ, ETH-Hönggerberg, 8093, Zurich²

and

Landau Institute for Theoretical Physics, Moscow, Russia³

In 2D XY ferromagnet the dipole force induces a strong interaction between spin-waves in the long-wavelength limit. The major effect of this interaction is the transformation of a propagating spin-wave into a diffusion mode. We study the anomalous dynamics of such diffusion modes. We find that the Janssen-De Dominicis functional, which governs this dynamics, approaches the non-Gaussian fixed-point. A spin-wave propagates by an anomalous anisotropic diffusion with the dispersion relation: $i\omega \sim k_y^{\Delta_y}$ and $i\omega \sim k_x^{\Delta_x}$, where $\Delta_y = 47/27$ and $\Delta_x = 47/36$. The low-frequency response to the external magnetic field is found.

I. INTRODUCTION

The notion of the ground state and independent elementary excitations lies in the background of the condensed matter theory. The excitations, such as electrons, holes, phonons, excitons, spin-waves etc. normally have a propagating, wave-like nature. In a homogeneous medium they are characterized by their momentum or quasi-momentum $\vec{\mathbf{p}}$ and their energy ω . Each kind of excitations has its specific dispersion relation or spectrum $\omega = \epsilon(\mathbf{p})$. A wide scope of physical problems can be solved assuming the excitation to be independent, or considering their interaction as a weak perturbation¹. However, in recent years more and more problems have occurred to go beyond the simple picture of non-interacting or weakly interacting excitations. In his pioneering work A.B.Migdal² has indicated that electron-phonon interaction is not weak in a narrow range of energy, leading to a strong renormalization of the Fermi-velocity and even to an instability of the Fermi-surface.

Recently a growing number of physical systems revealed excitations which bare spectrum, obtained from linearized equations of motion, is strongly distorted by interaction with vacuum and thermal fluctuations. A few recent examples are: the so-called marginal Fermi-liquid, in which electrons interacts via transverse magnetic field fluctuations³, and $\nu = 1/2$ state of the Fractional Quantum Hall effect in which initial electrons transform into quite different fermions⁴.

On the other hand, many years ago, the mode-mode interaction has been recognized as a necessary element of the critical dynamics^{5,6,7}. Particles, heat and spin diffusion must be considered as hydrodynamic modes in the long-wavelength limit, as well as propagating waves, such as sounds, spin waves etc. Their interaction has been proven to be substantial not only in the critical region, but also for the hydrodynamics of liquid crystals^{8,9} and for the CDW phason modes interacting with impurities¹⁰. The excitation spectrum of these systems is reconstructed by effects of strong interaction.

Here we present a new solvable and experimentally feasible situation where the strong interaction between spin-waves leads to the replacement of the propagating spin-wave by a

diffusion mode and to the appearance of a new soft-mode in a certain range of momentum. This is the two-dimensional XY ferromagnet with dipolar interaction between spins.

The spin-diffusion mode appears naturally in the paramagnetic phase and in the vicinity of the Curie point¹¹. We consider, on the other hand, a low temperature ordered phase, where no diffusion is expected but rather a propagating and weakly dissipating spin-wave mode.

In a 2D XY ferromagnet at low temperatures the dipolar interaction is relevant in the long wavelength limit, even despite of the low density of spin-waves. It was shown by one of the authors¹² that dipolar force induces an anomalous anisotropic scaling of spin-spin correlations in the ordered phase. In this article we find an analogous dynamical scaling.

This paper is organized as follows. In the next section we define the model and describe the spin-wave spectrum. In section III we discuss the dynamics of XY-magnet with the dipole interaction and formulate the perturbation expansion for the model, using the Janssen-De Dominicis¹³ technique. Section IV is devoted to the solution of the Dyson equation. There we find the self-induced dissipation of the spin-waves. In section V the renormalization of the diffusion mode is considered and the anomalous anisotropic exponents are found. In section VI the dynamical susceptibility is found. In conclusion we discuss prospects of the experimental observation of the anomalous modes. In the Appendix A we use the Ward-Takahashi identities to prove that the vertex corrections are small. Details of the Dyson equation solution can be found in the Appendixes B and C.

A brief report on main results of this article has been published earlier¹⁴.

II. THE HAMILTONIAN AND THE SPIN-WAVE SPECTRUM.

Spin-waves are fundamental excitations of the exchange magnet with spontaneously broken continuous symmetry. According to the so-called Adler principle¹⁵, the interaction between spin waves vanishes in the long-wavelength limit. At low temperatures the equilibrium density of spin waves is relatively small, and a long wavelength non-equilibrium

spin wave, excited by an external source, decays into other spin-waves or scatters on an equilibrium spin wave slowly. Thus, the dynamical properties of the exchange magnet are determined by the well-defined spin-wave mode. It also means that the imaginary part of the poles of the dynamical response function becomes much smaller than their real part as the wavelength grows to infinity.

The 2D exchange magnet with the easy-plane anisotropy is described by a following classical Hamiltonian:

$$H_{ExA}[\mathbf{S}] = \int d^2x \left[\frac{J}{2} (\nabla \mathbf{S})^2 + \frac{\lambda}{2} S_z^2 \right] - g_G \mu_B \int d^2x \mathbf{S} \mathbf{H}, \quad (1)$$

where J is the exchange coupling constant, λ is the strength of the easy-plane anisotropy ($\lambda > 0$)¹⁶, g is the dipole interaction coupling constant, $\mathbf{H}$ is the external magnetic field, g_G is gyromagnetic ratio, μ_B is the Bohr magneton and the field $\mathbf{S}(\mathbf{x})$ represents the local spin of the magnet and can be normalized by a constraint

$$\mathbf{S}^2(\mathbf{x}) = 1. \quad (2)$$

The magnetic dipolar energy H_{dip} is represented by the sum:

$$H_{dip} = \frac{g}{4\pi} \sum_{\mathbf{x}_i \neq \mathbf{x}_j} \frac{(\mathbf{S}_i \cdot \mathbf{S}_j) - 3(\mathbf{S}_i \cdot \hat{\nu})(\mathbf{S}_j \cdot \hat{\nu})}{|\mathbf{x}_i - \mathbf{x}_j|^3} \quad (3)$$

where $\hat{\nu}$ is a unit vector pointing from $\mathbf{x}_i$ to $\mathbf{x}_j$, $\mathbf{S}_i = \mathbf{S}(\mathbf{x}_i)$, and $g = 2\pi(g_G \mu_B S a^{-2})^2$ for the square lattice; for other lattices $g = 2\pi(g_G \mu_B S \sigma)^2$, where σ is the inverse area of a plaquette of the lattice. The magnetic dipole energy can be separated into a short-range and a long-range part in the standard way¹⁷. The short-range part renormalizes the single-ion spin anisotropy and favors in-plane spin orientation. The long-range part of the magnetic dipole energy is conveniently expressed in terms of the Fourier-transform $\mathbf{S}_{\mathbf{k}}$ of the local magnetization field $\mathbf{S}(\mathbf{x})$,

$$H_{dip} = \frac{g}{2} \sum_{\mathbf{k}} \frac{(\mathbf{S}_{\mathbf{k}} \mathbf{k})(\mathbf{S}_{-\mathbf{k}} \mathbf{k})}{|\mathbf{k}|}, \quad (4)$$

Thus, the total magnetic Hamiltonian is:

$$H = H_{ExA} + H_{dip}. \quad (5)$$

The dipole forces are crucial in the long-wavelength limit. Under the scaling transformation the short-range exchange interaction scales like L^{d-2} , whereas the dipole force scales like L^{2d-3} , where $d = 2, 3$ is the spatial dimension of a magnet and L is the scale. The dipole energy is characterized by the dipole constant g and the exchange energy is characterized by the exchange constant J . Normally in ferromagnets ga^{d-1} is much smaller than J (a is the lattice constant). However, as the scaling transformation shows, beyond the characteristic scale $L_d \sim (J/g)$ in 2D and $L_d \sim \sqrt{J/g}$ in 3D the dipole interaction dominates the energy of a spin-wave.

In the 2D XY ferromagnet the dipole force stabilizes the long-range order¹⁸. According to the Landau-Peierls-Hohenberg-Mermin-Wagner theorem (see e.g.¹⁹) in the absence of a long-range interaction the 2D magnet with broken continuous symmetry exhibits the algebraic decay of the spin correlations instead of the long-range order, and the infinite susceptibility at zero magnetic field. Thus, the dipole force plays a special role in 2D XY ferromagnet. In addition, the dipolar force is crucial for the spin statics and dynamics in the ordered phase. In contrast to the 3D ferromagnet, where the spin waves are almost free, in 2D the interaction between spin waves induced by dipole force is dominant in the long-wavelength limit.

The Hamiltonian (5) has two different scales: the anisotropy scale $L_A = \sqrt{J/\lambda}$ and the dipole length $L_D = J/g$. We assume that the anisotropy λ is large compared to the dipolar energy ($L_A < L_D$). We direct the y -axis along the net magnetization of the magnet and the z -axis perpendicular to the plane. The unit vector field $\mathbf{S}$ can be represented by two scalar fields $\phi(\mathbf{x}, t)$ and $\pi(\mathbf{x}, t)$

$$\mathbf{S} = \left(-\sqrt{1 - \pi^2} \sin \phi; \sqrt{1 - \pi^2} \cos \phi; \pi \right), \quad (6)$$

where both π and ϕ are small due to the fact that the dipole force stabilizes the long-range order.

With the precision to the fourth power of ϕ and π the Hamiltonian (5) is:

$$H[\phi] = \frac{1}{2} \int_{\mathbf{k}} \int_{\omega} \left((J\mathbf{k}^2 + h) \phi_{\mathbf{k},\omega} \phi_{-\mathbf{k},-\omega} + \lambda \pi_{\mathbf{k},\omega} \pi_{-\mathbf{k},-\omega} \right. \\ \left. + g \frac{\left(k_x (\phi_{\mathbf{k},\omega} - \phi_{\mathbf{k},\omega}^3/6) + k_y \left[\frac{\phi^2}{2} \right]_{\mathbf{k},\omega} \right) \left(k_x (\phi_{-\mathbf{k},-\omega} - \phi_{-\mathbf{k},-\omega}^3/6) + k_y \left[\frac{\phi^2}{2} \right]_{-\mathbf{k},-\omega} \right)}{|\mathbf{k}|} \right). \quad (7)$$

Here $\phi_{\mathbf{k},\omega}$ and $\pi_{\mathbf{k},\omega}$ are the Fourier-transforms of the fields $\phi(\mathbf{x}, t)$ and $\pi(\mathbf{x}, t)$ respectively. We expanded the in-plane magnetization components $\cos\phi$ and $\sin\phi$ up to the fourth power in small spin fluctuations ϕ . We take the uniform magnetic field $\mathbf{H}$ to be directed along the y -axis and $h = g_G \mu_B S H$. The Fourier-transformed quantities are defined by

$$\phi(\mathbf{x}, t) = \int_{\mathbf{k}} \int_{\omega} \phi_{\mathbf{k},\omega} e^{i(\mathbf{k}\mathbf{x} - \omega t)}, \quad (8)$$

where an abbreviated notation

$$\int_{\mathbf{k}} \int_{\omega} \equiv \iint \frac{d^2k}{(2\pi)^2} \frac{d\omega}{2\pi}$$

is used and $\left[\frac{\phi^2}{2} \right]_{\mathbf{k},\omega}$ denotes the Fourier-transformation of $\frac{\phi^2(\mathbf{x}, t)}{2}$.

The interaction between the spin waves is described by the non-quadratic part of the Hamiltonian (7):

$$H_{\text{int}} = \int_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3} \int_{\omega_1, \omega_2, \omega_3} f(\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3) \phi_{\mathbf{k}_1, \omega_1} \phi_{\mathbf{k}_2, \omega_2} \phi_{\mathbf{k}_3, \omega_3} \\ \times \delta(\mathbf{k}_1 + \mathbf{k}_2 + \mathbf{k}_3) \delta(\omega_1 + \omega_2 + \omega_3) \\ + \int_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4} \int_{\omega_1, \omega_2, \omega_3, \omega_4} g(\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4) \phi_{\mathbf{k}_1, \omega_1} \phi_{\mathbf{k}_2, \omega_2} \phi_{\mathbf{k}_3, \omega_3} \phi_{\mathbf{k}_4, \omega_4} \\ \times \delta(\mathbf{k}_1 + \mathbf{k}_2 + \mathbf{k}_3 + \mathbf{k}_4) \delta(\omega_1 + \omega_2 + \omega_3 + \omega_4), \quad (9)$$

where the three-leg bare vertex $f(\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3)$ is:

$$f(\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3) = \frac{g}{3} \sum_{i=1}^3 \frac{k_{ix} k_{iy}}{|\mathbf{k}_i|} \quad (10)$$

and the four-leg vertex is defined as:

$$g(\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4) = \frac{g}{24} \sum_{i>j=1}^4 \frac{(k_{iy} + k_{jy})^2 - (k_{ix} + k_{jx})^2}{|\mathbf{k}_i + \mathbf{k}_j|}. \quad (11)$$

The vertices (10) and (11) decrease as the momentum $\mathbf{k}$ goes to zero. Besides of that, these vertices are singular at $|\mathbf{k}| \rightarrow 0$. Nevertheless, the interaction between spin waves

asymptotically vanishes in the long wavelength limit for a 3D magnet¹⁵. No renormalization of the bare correlator

$$K(\mathbf{k}) = \langle \delta S_x(\mathbf{k}) \delta S_x(-\mathbf{k}) \rangle = \langle \phi_{\mathbf{k}} \phi_{-\mathbf{k}} \rangle = \frac{T}{J\mathbf{k}^2 + g \frac{k_x^2}{|\mathbf{k}|}} \quad (12)$$

appears in this limit (we put $h = 0$).

In 2D the situation changes drastically: the interaction grows with the wavelength, resulting in strong renormalization of critical exponents. To show the difference we calculate the upper marginal dimension of the Hamiltonian (7). Let us consider an arbitrary diagram from the perturbation expansion of some correlator. In order to add an internal line to such a diagram, we need to add three bare correlators K given by eq. (12), two vertices f and one integration over $dk_x d^{(D-1)}k_y$. From eq. (12) we see that for small momenta, $k_x \sim k_y^{3/2}$ and $K \sim k_y^{-2}$. Equation (10) gives $f \sim k_x \sim k_y^{3/2}$. Hence, if we require that the one-line insertion be dimensionless, we obtain: $3 \cdot (-2) + 2 \cdot 3/2 + 3/2 + (D-1) = 0$ or $D = 5/2$. It means that the theory is renormalizable in $D \leq 5/2$. It turns out that in the static 2D case the critical exponents can be found exactly.

Following the work¹², we rescale the field $\phi^2 \rightarrow \phi^2 / \sqrt{Jg}$ and rewrite the Hamiltonian (7) in a slightly different form:

$$H = \int_{\mathbf{k}} \left(a k_y^2 \phi_{\mathbf{k}} \phi_{-\mathbf{k}} + a^{-1} \frac{k_x^2}{|k_y|} \phi_{\mathbf{k}} \phi_{-\mathbf{k}} + 2w \frac{k_x k_y}{|k_y|} \phi_{\mathbf{k}} \left[\frac{\phi^2}{2} \right]_{-\mathbf{k}} + w^2 a |k_y| \left[\frac{\phi^2}{2} \right]_{\mathbf{k}} \left[\frac{\phi^2}{2} \right]_{-\mathbf{k}} \right), \quad (13)$$

where $a = \sqrt{J/g}$, $w^2 = \sqrt{g/J^3}$ and we have taken into account that for small $\mathbf{k}$ one can substitute $|\mathbf{k}| \rightarrow k_y$ (see (12)). Requiring that the Hamiltonian (13) does not change under scale transformation $k_y \rightarrow l k_y$, $k_x \rightarrow l^{\Delta_x^0} k_x$, $\phi_{\mathbf{k}} \rightarrow l^{\Delta_\phi^0} \phi_{\mathbf{k}}$, $a \rightarrow l^{\Delta_a^0} a$, $w \rightarrow l^{\Delta_w^0} w$, we find the bare exponents $\Delta_x^0 = 3/2$, $\Delta_a^0 = 0$, $\Delta_\phi^0 = -9/4$, $\Delta_w^0 = 1/4$ (where we have taken into account that $\left[\frac{\phi^2}{2} \right]$ has one more integration). Now, according to the standard procedure²⁰, we introduce the renormalized field and charges: $\phi^2 = Z_\phi \phi_R^2$, $a = Z_a a_R$, $w = Z_w \mu^{1/4} \tilde{w}_R$, where μ is the scale at which the dimensionless $\tilde{w}_R$ is observed. The second term in the Hamiltonian (13) is non-analytic at small momenta. It means that there can be no corrections to this term in the regular perturbation expansion. Furthermore, from Appendix A one can see,

that in this case, due to symmetry, there will be no correction to the third term either. Hence, we can establish the relations between the renormalization coefficients:

$$Z_a^{-1} Z_\phi = 1 \quad (14)$$

$$Z_\phi^{3/2} Z_w \mu^{1/4} = 1. \quad (15)$$

It can be proven that the renormalization group flow has a non-zero fixed point. It means that $\tilde{w}_R = \text{const}$ and hence $Z_w \sim (\Lambda/\mu)^{1/4}$, where Λ is the scale at which the coupling constant equals w . Introducing the critical exponents $Z_\phi \sim \Lambda^{2\Delta_\phi}$, $Z_a \sim \Lambda^{\Delta_a}$, we find from (14) and (15) that $\Delta_\phi = -1/12$ and $\Delta_a = -1/6$. Demanding that the first and the second terms in the Hamiltonian (13) have the same dimensions, we find $\Delta_x = 4/3$. Finally, from the Callan-Symanzik equation²⁰, the long range limit of the two point correlation function can be found:

$$K(\mathbf{x}) = (x^2 + |y|^{8/3})^{-1/4} f(x/|y|^{4/3}),$$

where $f(x)$ is an arbitrary finite function. The dynamical properties of the magnet, however, are still to be found.

III. THE DYNAMICS OF XY MAGNET

At $T = 0$ the classical magnet with the Hamiltonian (5), obeying the constraint (2), follows the Landau-Lifshitz equations (see e.g.²¹):

$$S\hbar \frac{\partial \mathbf{S}(\mathbf{x}, t)}{\partial t} = \mathbf{S}(\mathbf{x}, t) \times \frac{\delta H}{\delta \mathbf{S}(\mathbf{x}, t)}, \quad (16)$$

where S is the absolute value of a spin localized on a magnetic ion. In terms of the canonically conjugated fields π and ϕ , eq. (16) can be rewritten in the Hamiltonian form:

$$\hbar S \partial_t \pi(\mathbf{x}, t) = \frac{\delta H}{\delta \phi(\mathbf{x}, t)}, \quad (17)$$

$$-\hbar S \partial_t \phi(\mathbf{x}, t) = \frac{\delta H}{\delta \pi(\mathbf{x}, t)} \approx \lambda \pi(\mathbf{x}, t). \quad (18)$$

In the harmonic approximation equations (17,18) imply the dispersion relation for the spin-wave mode¹⁶ in a XY magnet:

$$\epsilon^2(\mathbf{k}) = \lambda \left(J\mathbf{k}^2 + g \frac{k_x^2}{|\mathbf{k}|} \right) = c^2 \left(\mathbf{k}^2 + p_0 \frac{k_x^2}{|\mathbf{k}|} \right), \quad (19)$$

where $c = \sqrt{\lambda J}$ is the spin-wave velocity and $p_0 = g/J$. The out-of-plane anisotropy λ affects the dynamics in the long-wavelength limit $k \ll p_\lambda = \sqrt{\lambda/J}$ considered in this article. Thus, p_λ is the upper cut-off momentum in our theory. A spin wave with $k \gg p_0$ has the phonon-like isotropic spectrum $\epsilon = ck$. The range $p_0 \ll k \ll p_\lambda$ will be called acoustic shell ($\mathcal{A}$ -shell). At lower momenta $p \ll p_0$ the spin-wave spectrum is dominated by the dipolar interaction: $\epsilon(\mathbf{k}) \approx c\sqrt{p_0 k} \sin \theta$, where θ is the angle between the direction of the spontaneous magnetization and the wave-vector. This range of momenta will be called dipolar shell ($\mathcal{D}$ -shell). The effect of the presence of the dipolar force in the 2D XY-magnet is not limited to the change of free spin-wave spectrum. As we have mentioned earlier, it leads to strong spin-wave interaction and to a crucial transformation of the spin propagation.

Without dipolar forces the dynamics of the 2D XY ferromagnet is well described by non-interacting spin waves. The high level of fluctuations leads to a strong temperature dependence of the dynamic spin correlators, which have algebraic character, just as static ones^{22,23}. The renormalized two-point spin correlation function features the pole with the temperature dependent power exponent. The dipole force suppresses such strong spin fluctuations, but not entirely.

In addition, the dipolar interaction induces decay processes. As a result finite spin-wave life-time $\Gamma(\mathbf{k})$ or the width of the level $b(\mathbf{k}) = \Gamma^{-1}(\mathbf{k})$ appears. In 3D at low temperature and at small momentum $|\mathbf{k}|$, the width $b(\mathbf{k})$ is much smaller than $\omega(\mathbf{k})$. In 2D, however, the interaction is essential and must be considered seriously.

To take into account the dissipation induced by thermal fluctuations at a temperature T , we introduce a phenomenological dissipation functional²⁴:

$$R[\phi] = \int dt d^2x d^2x' R(\mathbf{x} - \mathbf{x}') \dot{\phi}(\mathbf{x}) \dot{\phi}(\mathbf{x}'). \quad (20)$$

Eliminating π from equations (17,18) and adding a proper dissipation term, one obtains a following equation for $\phi(\mathbf{x}, t)$:

$$-\frac{1}{\lambda}\partial_t^2\phi(\mathbf{x}, t) = \left[\frac{1}{\lambda}\epsilon^2(\mathbf{k})\phi(\omega, \mathbf{k})\right]_{\mathbf{x}, t} + \frac{\delta H_{\text{int}}}{\delta\phi(\mathbf{x}, t)} + \frac{\delta R}{\delta\dot{\phi}(\mathbf{x}, t)} + \eta(\mathbf{x}, t) - h(\mathbf{x}, t), \quad (21)$$

where $h(\mathbf{x}, t)$ is the external magnetic field, and the interaction part of the Hamiltonian is determined by equation (9). We have introduced the random noise $\eta(\mathbf{x}, t)$ in equation (21). The noise, in effect, generates dissipation. As usual, the random noise is assumed to obey the Gaussian statistics. Its correlation function is determined by the fluctuation-dissipation theorem²⁵:

$$\langle\eta_{\mathbf{k}}\eta_{-\mathbf{k}}\rangle = 2TR(\mathbf{k}) \quad (22)$$

Here $R(\mathbf{k})$ is the Fourier-transform of the function $R(\mathbf{x}-\mathbf{x}')$. The dissipation in the exchange ferromagnet vanishes in the long-wavelength limit¹⁵: $R(\mathbf{k}) = b\mathbf{k}^2$. In 2D XY dipole magnet the dissipation does not vanish in the long-wavelength limit. We notify:

$$R(\mathbf{k}) = \frac{1}{\Gamma(\mathbf{k})} \quad (23)$$

We emphasize that the finite life-time Γ is determined self-consistently by the processes of the decay and scattering of spin waves. We neglect the spin-wave-electron and spin-wave-sound interactions. The first interaction is not weak, but the Fermi-velocity is much higher than the spin-wave velocity and the spin-electron interaction is not effective for long wavelength. Even the sound velocity can be much larger than the spin-wave velocity, since the latter is proportional to small $\sqrt{\lambda}$.

Our aim then is to calculate the linear response function $G(\mathbf{x}, t)$ to a weak external field $h(\mathbf{x}, t)$ (see equation (21)) averaged over the thermal fluctuations $\eta(\mathbf{x}, t)$. We apply the Janssen-De Dominicis functional method¹³ to reformulate stochastic equation (21) in terms of the path integral. The probability distribution for the noise $\eta(\mathbf{x}, t)$ is:

$$W[\eta] \sim \exp\left[-\frac{1}{4T}\int d^2x d^2x' \int dt \eta(\mathbf{x}, t) R^{-1}(\mathbf{x} - \mathbf{x}') \eta(\mathbf{x}', t)\right]. \quad (24)$$

Following the standard dynamic field theory^{20,9}, upon averaging over the noise distribution and introducing auxiliary response field $\hat{\phi}(\mathbf{x}, t)$, one can reduce the solution of the stochastic differential equation (21) to the calculation of the dynamical partition function:

$$Z[j, \hat{j}] = \int \mathcal{D}[\phi] \mathcal{D}[i\hat{\phi}] \exp\left(\mathcal{J}[\phi, \hat{\phi}] + \int d^2x \int dt [\hat{j}\hat{\phi} + j\phi]\right) \quad (25)$$

and its derivatives over the currents j and $\hat{j}$. Here $\mathcal{J}(\phi, \hat{\phi})$ is the Janssen-De Dominicis functional (JDF).

$$\begin{aligned} \mathcal{J}[\phi, \hat{\phi}] = & \int d^2x \int d^2x' \int dt \hat{\phi}(\mathbf{x}, t) T R(\mathbf{x} - \mathbf{x}') \hat{\phi}(\mathbf{x}', t) \\ & - \hat{\phi}(\mathbf{x}', t) \left[\frac{1}{\lambda} \partial_t^2 \phi(\mathbf{x}, t) \delta(\mathbf{x}' - \mathbf{x}) + \frac{\delta H}{\delta \phi(\mathbf{x}, t)} \delta(\mathbf{x}' - \mathbf{x}) + R(\mathbf{x}' - \mathbf{x}) \partial_t \phi(\mathbf{x}, t) \right]. \end{aligned} \quad (26)$$

By differentiation of the JDF over j and $\hat{j}$ one can obtain any correlation function.

We define the inverse bare propagator $\hat{\mathbf{G}}_0^{-1}$

$$\hat{\mathbf{G}}_0^{-1} = \begin{pmatrix} -2TR(\mathbf{k}) & \frac{\omega^2}{\lambda} - \frac{\epsilon^2(\mathbf{k})}{\lambda} - i\omega R(\mathbf{k}) \\ \frac{\omega^2}{\lambda} - \frac{\epsilon^2(\mathbf{k})}{\lambda} + i\omega R(\mathbf{k}) & 0 \end{pmatrix}, \quad (27)$$

so that the harmonic ("Gaussian") part of $\mathcal{J}$ can be written in the form:

$$\mathcal{J}_0[\phi, \hat{\phi}] = -\frac{1}{2} \int_{\mathbf{k}} \int_{\omega} \begin{pmatrix} \hat{\phi}(\omega, \mathbf{k}) & \phi(\omega, \mathbf{k}) \end{pmatrix} \hat{\mathbf{G}}_0^{-1}(\omega, \mathbf{k}) \begin{pmatrix} \hat{\phi}(-\omega, -\mathbf{k}) \\ \phi(-\omega, -\mathbf{k}) \end{pmatrix}. \quad (28)$$

From eq. (27) one finds the bare propagator:

$$\hat{\mathbf{G}}_0 = \begin{pmatrix} 0 & G_0^*(\omega, \mathbf{k}) \\ G_0(\omega, \mathbf{k}) & D_0(\omega, \mathbf{k}) \end{pmatrix}, \quad (29)$$

where we define the bare dynamical response and spin-spin correlation function as follows:

$$G_0(\omega, \mathbf{k}) = \frac{\lambda}{\omega^2 - \epsilon^2(\mathbf{k}) - i\omega\lambda R(\mathbf{k})}, \quad (30)$$

$$D_0(\omega, \mathbf{k}) = \frac{2T\lambda R(\mathbf{k})}{[\omega^2 - \epsilon^2(\mathbf{k})]^2 + \omega^2\lambda^2 R^2(\mathbf{k})}. \quad (31)$$

They obey the standard fluctuation-dissipation relation: $D_0 = \frac{2T}{\omega} \text{Im} G_0$. The same relation is correct for the total dynamic correlation $D(\omega, \mathbf{k})$ and the total linear response function $G(\omega, \mathbf{k})$: $D = \frac{2T}{\omega} \text{Im} G$.

The anharmonic (interaction) part of $\mathcal{J}$ is:

$$\begin{aligned} \mathcal{J}_{\text{int}} = & \sum_{\omega, \mathbf{k}} \hat{\phi}_{\omega, \mathbf{k}} \frac{\delta H_{\text{int}}}{\delta \phi_{\omega, \mathbf{k}}} = \sum_{\omega, \mathbf{k}} \hat{\phi}_{\omega, \mathbf{k}} \left(3 \sum_{\omega_2, \omega_3} \sum_{\mathbf{k}_2, \mathbf{k}_3} f(\mathbf{k}, \mathbf{k}_2, \mathbf{k}_3) \phi_{\mathbf{k}_2, \omega_2} \phi_{\mathbf{k}_3, \omega_3} \right. \\ & \times \delta(\mathbf{k} + \mathbf{k}_2 + \mathbf{k}_3) \delta(\omega + \omega_2 + \omega_3) + 4 \sum_{\omega_2, \omega_3, \omega_4} \sum_{\mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4} g(\mathbf{k}, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4) \phi_{\mathbf{k}_2, \omega_2} \phi_{\mathbf{k}_3, \omega_3} \phi_{\mathbf{k}_4, \omega_4} \\ & \left. \times \delta(\mathbf{k} + \mathbf{k}_2 + \mathbf{k}_3 + \mathbf{k}_4) \delta(\omega + \omega_2 + \omega_3 + \omega_4) \right), \end{aligned} \quad (32)$$

We define in a common way the self-energy operator $\hat{\Sigma}(\omega, \mathbf{k})$ by the relation:

$$\hat{\mathbf{G}}^{-1}(\omega, \mathbf{p}) = \hat{\mathbf{G}}_0^{-1}(\omega, \mathbf{p}) - \hat{\Sigma}(\omega, \mathbf{p}), \quad (33)$$

where:

$$\hat{\mathbf{G}} = \begin{pmatrix} 0 & G^*(\omega, \mathbf{k}) \\ G(\omega, \mathbf{k}) & D(\omega, \mathbf{k}) \end{pmatrix} \quad (34)$$

and $G(\omega, \mathbf{k})$ and $D(\omega, \mathbf{k})$ are the complete response function and correlator respectively.

The self-energy $\hat{\Sigma}(\omega, \mathbf{k})$ satisfies Dyson equation:

$$\hat{\Sigma}(\omega, \mathbf{k}) = g \int_{\Omega} \int_{\mathbf{p}} \hat{\mathbf{G}}(\Omega, \mathbf{p}) \hat{\Lambda}(\mathbf{p}, \mathbf{k}; \omega, \Omega) \hat{\mathbf{G}}(\omega - \Omega, \mathbf{k} - \mathbf{p}), \quad (35)$$

where $\hat{\Lambda}(\mathbf{p}, \mathbf{k}; \omega, \Omega)$ is the full vertex. The bare vertex is given by equation:

$$\hat{\Lambda}_0(\mathbf{p}, \mathbf{k}; \omega, \Omega) = \frac{2TR(\mathbf{p})}{\lambda} f^2(\mathbf{p}, \mathbf{k}) \begin{pmatrix} -D_0(\Omega, \mathbf{p}) & G_0^*(\Omega, \mathbf{p}) \\ G_0(\Omega, \mathbf{p}) & 0 \end{pmatrix}. \quad (36)$$

We have denoted $f(\mathbf{p}, \mathbf{k}) \equiv f(\mathbf{p}, \mathbf{k}, \mathbf{p} - \mathbf{k})$.

Further we consider a limit $R(\mathbf{k}) \rightarrow +0$. According to the FDT, the matrix $\hat{\Sigma}(\omega, \mathbf{p})$ must have a form:

$$\hat{\Sigma}(\omega, \mathbf{p}) = \begin{pmatrix} \frac{2T}{\lambda\omega} \text{Im}\Sigma(\omega, \mathbf{p}) & \frac{1}{\lambda} \Sigma^*(\omega, \mathbf{p}) \\ \frac{1}{\lambda} \Sigma(\omega, \mathbf{p}) & 0 \end{pmatrix}, \quad (37)$$

where the self-energy function $\Sigma(\omega, \mathbf{p})$ is associated with the complete response function $G(\omega, \mathbf{p})$ by the same relationship:

$$G^{-1}(\omega, \mathbf{p}) = G_0^{-1}(\omega, \mathbf{p}) - \frac{1}{\lambda} \Sigma(\omega, \mathbf{p}). \quad (38)$$

In the Appendix A we show that the full vertex Λ can be approximated with a good accuracy by its bare value $f(\omega, \mathbf{p})$ in the low-frequency range. In this respect our theory reminds the Migdal theory of the interacting electron-phonon system². In the Migdal theory the simplification is due to a narrow scale of the energy shell in which the interaction proceeds. In our theory we assume that the frequency of spin fluctuations is small instead.

We notify the real and the imaginary part of the self-energy term as: $\Sigma = a^2(\omega, \mathbf{p}) - i\omega b(\omega, \mathbf{p})$. Thus, the Green function (33) reads:

$$G^{-1}(\omega, \mathbf{p}) = \omega^2 - \epsilon^2(\mathbf{p}) - a^2(\omega, \mathbf{p}) + i\omega b(\omega, \mathbf{p}), \quad (39)$$

while the spin-spin correlation function is:

$$D(\omega, \mathbf{p}) = \frac{b(\omega, \mathbf{p})}{[\omega^2 - \epsilon^2(\mathbf{p}) - a^2(\omega, \mathbf{p})]^2 + \omega^2 b^2(\omega, \mathbf{p})} \quad (40)$$

(we have slightly changed the definitions of G and D and referred the factor $\sqrt{2T\lambda^3}$ to the vertex).

We employ the reduced temperature $t = T/4\pi J$ and the ratio $g/\sqrt{J\lambda} = p_0/\sqrt{\lambda/J}$ as small parameters. The latter means that the $\mathcal{A}$ -shell is much larger than the $\mathcal{D}$ -shell. We also use the notation $L = \log(\sqrt{J\lambda}/g)$.

The main contribution to the self-energy is given by the one-loop diagrams shown in Fig.1a, 1b. Our theory is valid only if the temperature is small:

$$t \log(\sqrt{J\lambda}/g) = tL \ll 1. \quad (41)$$

Under this condition the two-loop corrections are small, and the diagram Fig.1b contributes to a negligible change of the spectrum (19)²⁰. Such neglecting of the two-loop diagrams (vertex correction) was a major assumption in the so-called mode-coupling methods¹¹. This approximation serves well in the theory of the 3D critical dynamics with the dipole force being included. Later we prove this assumption for 2D. Thus, the Dyson equation for our problem is as follows:

$$\Sigma(\Omega, \mathbf{q}) = 18\lambda^3 T \int_{\mathbf{p}} \int_{\omega} f^2(\mathbf{p}, \mathbf{q}) D(\omega, \mathbf{p}) G(\omega + \Omega, \mathbf{q} - \mathbf{p}), \quad (42)$$

The functions $b(\omega, \mathbf{p})$ and $a(\omega, \mathbf{p})$ are even in both arguments¹. The imaginary part of the self-energy is odd in ω : $\text{Im}\Sigma(\Omega, \mathbf{q}) = -\Omega b(\Omega, \mathbf{q})$. Hence, equation for the dissipation function reads:

$$b(\Omega, \mathbf{q}) = 9\lambda^3 T \int_{\mathbf{p}} \int_{\omega} f^2(\mathbf{p}, \mathbf{q}) D(\omega, \mathbf{p}) D(\omega + \Omega, \mathbf{q} - \mathbf{p}). \quad (43)$$

The integrand in (43) is positive. Thus, the main contribution to $b(\Omega, \mathbf{q})$ comes from the region, where poles of the two D -functions coincide. The function $D(\omega, \mathbf{p})$ has poles at $\omega \approx \pm\epsilon(\mathbf{p})$ in the $\mathcal{A}$ -shell. Following the terminology of the field theory, we call the surface $\omega^2 = \epsilon^2(\mathbf{p})$ the mass-shell. The self-energy in the $\mathcal{A}$ -shell is small as it is shown in the Appendix B and we neglect it. Because the dissipation is small, the D -function can be represented as a sum of δ -functions:

$$D(\omega, \mathbf{p}) \approx \sum_{\pm} \frac{\pi}{2\epsilon^2(\mathbf{p})} \delta(\Delta\omega_{\pm}), \quad (44)$$

where $\Delta\omega_{\pm} = \omega \pm \epsilon(\mathbf{p})$ measures the deviation from the mass-shell. After integrating ω out from eq. (43) with the D -functions from (44), we recover the Fermi Golden Rule for the probability of the spin-wave decay and scattering processes.

Looking for the long wavelength quasi-excitations, we need the self-energy at very small momenta $q \ll p_0$, which we denote as Σ_0 . We expect the quasi-excitations to be soft: $\Omega \ll cq$. Here we restrict the quasi-excitation wave-vector $\mathbf{q}$ to be directed almost along the magnetization: $|q_x| \ll q$ (arbitrarily directed $\mathbf{q}$ are considered in Appendix C). The essential contribution to the integral in equation (42) comes from the internal momentum p being in the $\mathcal{A}$ -shell and the internal frequency $\omega = \epsilon(\mathbf{p})$. Performing the integration over ω with the D -function from equation (44), we find (for derivation see Appendix B and note that we neglect the real part of the self-energy in the $\mathcal{A}$ -shell, as it is justified in the Appendix B):

$$\Sigma_0 = \frac{c^2 p_0^2 t}{4\pi} \int \frac{c^4 p^3 dp}{\epsilon^4(\mathbf{p})} \frac{\Omega \sin^2(2\psi) d\psi}{\Omega - cq \cos \psi + ib_1}, \quad (45)$$

where b_1 is the dissipation function $b(\omega, \mathbf{p})$ of a spin-wave inside the $\mathcal{A}$ -shell. Note, that $\text{Re}\Sigma_0$ vanishes in the static limit $\Omega = 0$.

If $cq \gg b_1$, we make the integral over ψ in equation (45) to find:

$$\Sigma_0(\chi) = c^2 p_0^2 t L \cos^2 \chi \exp(-2i\chi), \quad (46)$$

where χ , defined by the equation $\cos \chi = \Omega/cq$, measures the deviation from the mass shell. More generally, we introduce a notation r for the ratio Ω/cq ($r = \cos \chi$ if $r \leq 1$). Then

$$\Sigma_0(r) = c^2 p_0^2 L t r^2 (2r^2 - 1 - 2r\sqrt{r^2 - 1}). \quad (47)$$

Note that $\Sigma_0(r) \approx -c^2 p_0^2 t/4$ when $r \rightarrow \infty$ and $|\Sigma_0(r)| < c^2 p_0^2 t/4$ at any r . The self-energy $\Sigma_0(r)$ is real for $r > 1$.

If q is so small that $cq \ll b_1$, eq. (45) implies the q -independent dissipation constant:

$$b_0 = c^2 p_0^2 t L \int \frac{d\psi}{4\pi} \frac{\sin^2(2\psi)}{b_1(\psi)}. \quad (48)$$

In this calculation we have used the fact that the dissipation of a spin-wave in the $\mathcal{A}$ -shell b_1 depends only on the angle ψ between the direction of magnetization and the spin-wave wave-vector $\mathbf{p}$ which we prove below.

Now we need to calculate $b_1(\psi)$. An unusual feature of our theory is that the dissipation process in the $\mathcal{A}$ -shell is mediated by an off-mass-shell virtual spin wave. Indeed, the dispersion relation (19) does not allow for decay or merging processes. Alternatively, as we will show, the dissipation of a spin-wave in the $\mathcal{A}$ -shell, propagating along the direction specified with the angle ψ ($\sin \psi = q_x/q$), is mediated by an internal virtual spin-wave in (9), with a momentum of $p \ll p_0$ and a frequency of $\omega < cp$, propagating along the direction very close to the y -axis $\varphi^2 \ll 1$ (where $\sin \varphi = p_x/p$), to provide a finite attenuation of this state. The integration over ω with one of the D -functions in (43), taken in the form (44), leads to a following equation:

$$b_1 = -9c^4 t \frac{f^2(\mathbf{0}, \mathbf{q})}{8\pi J^2 q^2} \int d^2\mathbf{p} D(\epsilon(\mathbf{p} + \mathbf{q}) - \epsilon(\mathbf{q}), \mathbf{p}). \quad (49)$$

Since $\omega = \epsilon(\mathbf{p} + \mathbf{q}) - \epsilon(\mathbf{q})$, we conclude that $\omega = cp \cos \Phi$, where $\Phi = \psi - \varphi \approx \psi$ is the angle between the vectors $\mathbf{q}$ and $\mathbf{p}$. According to fluctuation-dissipation theorem: $D(\epsilon_{\mathbf{p}+\mathbf{q}} -$

$\epsilon_{\mathbf{q}, \mathbf{p}} = \frac{1}{\epsilon_{\mathbf{p}+\mathbf{q}} - \epsilon_{\mathbf{q}}} \text{Im} G(\epsilon_{\mathbf{p}+\mathbf{q}} - \epsilon_{\mathbf{q}}, \mathbf{p})$. Invoking the definition of the angle χ for virtual spin-wave, we find that $\chi = \Phi \approx \psi$. Substituting $f^2(\mathbf{0}, \mathbf{q}) = (1/9) q^2 \sin^2(2\psi)$, $\epsilon^2(\mathbf{p}) = c^2 p^2 + c^2 p_0 p \sin^2 \varphi \approx c^2 p^2 + c^2 p_0 p \varphi^2$ and taking into account that Σ_0 from (46) depends only on $\chi = \psi$, we write:

$$b_1(\psi) = \frac{c^2 p_0^2 t}{2\pi} \text{Im} \int \frac{\sin^2 \psi \cos \psi \, dp d\varphi}{p^2 \sin^2 \psi + p_0 p \varphi^2 + \Sigma_0(\psi)/c^2}. \quad (50)$$

Note that the most dangerous region of integration is the region of small p , such that $\varphi^2 \sim p/p_0 \ll 1$ in (50). In other words, the dissipation of a short wavelength spin-wave, propagating in the direction ψ , is determined by the scattering on the long wavelength virtual spin-wave, with the momentum along $\hat{y}$ -direction, which lies on a specific distance off the mass-shell: $\omega/cp = \cos \psi$. The integration over p in (50) is confined towards the crossover region: $p \sim p_c = p_0 \sqrt{tL}$.

To find the anisotropic dissipation of a spin-wave mode in the $\mathcal{A}$ -shell, we plug $\Sigma_0(\psi)$ from equation (46) into (50). After a change of variables $(p, \varphi) \rightarrow (\rho, \vartheta)$, given by formulae $p = p_0 \rho^2 \cos \vartheta$ and $\varphi = \rho \sin \psi \sin \vartheta \cos^{-1/2} \vartheta$ ($-\infty < \rho < \infty$ and $0 < \vartheta < \pi/2$), the integration becomes trivial and gives:

$$b_1(\psi) = \beta_1 t^{3/4} c p_0 \frac{\sin^{3/2}(2\psi) \sin(\psi/2)}{L^{1/4} \cos \psi}, \quad (51)$$

where the direction of the spin-wave is limited to the fundamental quadrant: $0 < \psi < \pi/2$, and $\beta_1 = \Gamma^2(1/4)/4\sqrt{2\pi} \approx 1.31$.

Let us return to the range of very low momenta $p \ll b_1/c$. Plugging (51) into (48), one finds:

$$b_0 = \beta_0 c p_0 t^{1/4} L^{5/4},$$

where $\beta_0 \approx 1.24$. The condition $c p_{DM} \sim b_1$ defines the crossover wave-vector: $p_{DM} \sim \beta_1 p_0 t^{3/4} / L^{1/4}$, between the self-energies (46) and (48). The dissipation functions (46, 48 and 51) represent the self-consistent solution of the Dyson equation (42, 43).

Finally, we verify that the two-loop correction (see Fig.1c) is negligible. There exists several diagrams with different arrangements of G and D functions. On each of the two

short loops on the diagram Fig.1c there exists at least one D function but there may be two of them. We consider only the most 'dangerous' diagram with each short loop having exactly one D -function (see Fig.2). Note that the main contribution to the diagram Fig.2 comes from regions of internal momenta $\mathbf{p}_1$ and $\mathbf{p}_2$ are restricted to the $\mathcal{A}$ -shell. Inside the $\mathcal{A}$ -shell the Green and the D -functions have strong singularities on the mass-shell. As it was done in the Appendix B we integrate in both short loop the internal frequencies ω_1 and ω_2 and find that only non-static term is non-zero:

$$\Sigma_0 \sim t^2 \int \frac{d^2 p_1 d^2 p_2}{\epsilon^3(\mathbf{p}_1) \epsilon^3(\mathbf{p}_2)} \frac{\Omega f^2(\mathbf{p}_1, \mathbf{p}_2) f(\mathbf{p}_1, \mathbf{p}_1) f(\mathbf{p}_2, \mathbf{p}_2)}{(\Omega - cq \cos \phi_1 - ib_1)(\Omega - cq \cos \phi_2 - ib_2)} \times \frac{\epsilon^{-2}(\mathbf{p}_1 - \mathbf{p}_2)}{(\epsilon(\mathbf{p}_1) - \epsilon(\mathbf{p}_2) - \epsilon(\mathbf{p}_1 - \mathbf{p}_2))}, \quad (52)$$

where ϕ_1 and ϕ_2 is the direction of the momenta $\mathbf{p}_1$ and $\mathbf{p}_2$. We assumed that $\phi_1 \approx \phi_2$. Since the three spin-wave processes is not allowed by the conservation laws, the Green function $G(\omega_1 - \omega_2, \mathbf{p}_1 - \mathbf{p}_2)$ is off the mass-shell (the corresponding last denominator in Eq.(52) reads: $(p_1 + p_2)(\phi_1 - \phi_2)^2 + p_0 \cos^2(\phi_1) > 0$ (if $p_1, p_2 \gg p_0$). Now we can count the momenta powers in Eq.(52) to verify that the integration is convergent towards the dipolar momentum p_0 , and, thus has no logarithm. A simple counting of temperatures shows that Eq.(52) $\sim t^2/b_1^2 \sim t^{1/2}$. Hence the two-loop dissipation function is $b'_0 = b_0(1 + t^{1/4}/L)$. Similar consideration shows that the function $b'_1(\psi, q) - b_1(\psi)$, which represents the two-loop corrections for b_1 , is small in $t^{1/4}$, and is also small in the ratio p_0/q .

Having explicit expressions for the self-energy we can analyze the dispersion relation $\omega^2 = \epsilon^2(\mathbf{p}) + \Sigma(\omega, \mathbf{p})$ in the range of small ω and p . New results are expected for the region $p < p_c = p_0 \sqrt{tL}$ in which Σ_0 becomes comparable with $\epsilon^2(\mathbf{p})$. In a range of momentum $p_{DM} \ll p \ll p_c$ and angles $\psi \ll \sqrt{p_0 t L / p}$, we find a new propagating soft mode with the dispersion:

$$\omega = cp(p^2 + p_0 p \psi^2)^{1/2} / p_0 \sqrt{tL}. \quad (53)$$

The dissipation of the soft mode grows to the boundary of the region and becomes of the order of its energy at $\psi \sim \sqrt{p_0 t L / p}$ or $p \sim p_{DM}$. There is no soft mode beyond the

indicated range. The spin-wave mode persists at $p > p_0 t L$. In a range $p \ll p_{DM}$ and small angles a new diffusion mode occurs with the dispersion:

$$\omega = -i\epsilon^2(\mathbf{p})t^{-1/4}L^{5/4}/\beta_0 c p_0. \quad (54)$$

The angular range of the diffusion mode increases with decreasing p and captures the entire circle at $p < p_0 t L$.

At the end of this section we would like to remind that equation (48) was obtained for the quasi-excitation directed along the y -axis. It can be easily checked that in case of arbitrarily directed $\mathbf{q}$ one must write $\sin^2(2\psi + 2\phi)$ instead of $\sin(2\psi)$ in eq. (45) where $\sin \phi = q_x/q$. However, in this case the integral becomes singular, so one must treat this equation more carefully. We will come back to this question in Appendix C.

IV. RENORMALIZATION OF THE DIFFUSION MODE.

In this section we concern ourself with the renormalization of the diffusion mode. As we established in the previous section, at wavevectors $p < p_{DM}$ the diffusive dynamics term dominates ($\lambda \rightarrow 0$ limit) in the harmonic part of JD Functional (27-28). The interaction between 'diffusons', given by the anharmonic part of the JDF (32), effectively 'renormalizes' the 'diffuson' dispersion at very small wavevectors $p < p_a \ll p_{DM}$. We shall determine the anomalous diffusion onset wavevector p_a in the end of this section.

To simplify further calculations, we introduce a scale transformation of the fields $\phi, \hat{\phi} \rightarrow (Jg/T^2)^{-1/4}\phi, (Jg/T^2)^{-1/4}\hat{\phi}$. In these notations the JDF (28,32) is:

$$\begin{aligned} \mathcal{J}[\phi, \hat{\phi}] = & \sum_{\omega, \mathbf{k}} \hat{\phi}_{-\omega, -\mathbf{k}} \left(a \frac{T}{\Gamma_0} \hat{\phi}_{\omega, \mathbf{k}} - \left(a k_y^2 + \frac{k_x^2}{a|k_y|} \right) \phi_{\omega, \mathbf{k}} - a \frac{i\omega}{\Gamma_0} \phi_{\omega, \mathbf{k}} \right. \\ & \left. - w \frac{k_x k_y}{|k_y|} \left[\frac{\phi^2}{2} \right]_{\omega, \mathbf{k}} - w \sum_{\Omega, \mathbf{p}} \phi_{\omega-\Omega, \mathbf{k}-\mathbf{p}} \frac{p_x p_y}{|p_y|} \phi_{\Omega, \mathbf{p}} - w^2 a \sum_{\Omega, \mathbf{p}} \phi_{\omega-\Omega, \mathbf{k}-\mathbf{p}} |p_y| \left[\frac{\phi^2}{2} \right]_{\Omega, \mathbf{p}} \right), \end{aligned} \quad (55)$$

where $\Gamma_0 = J\Gamma$, $a = \sqrt{J/g}$ and

$$w = (T^2 g / J^3)^{1/4}. \quad (56)$$

Note, that one can get rid off the spatial anisotropy charge a by rescaling the x coordinate only.

Assuming that all terms in the r.-h.s. of JDF (55) have the same scaling dimensions, equal to zero, and accepting the dimension of k_y $\Delta_y = 1$, $\Delta_a^0 = 0$ and $\Delta_{1/\Gamma}^0 = 0$, we immediately find from the second and third terms that $\Delta_{k_x}^0 = 3/2$. Comparing the second and the fourth terms, we get $\Delta_\omega^0 = 2$. The first and the fourth terms give $\tilde{\Delta}_\phi^0 + \Delta_\omega^0 = \tilde{\Delta}_\phi^0$. From the condition that the scaling dimension of the first term is zero, we have $2\tilde{\Delta}_\phi^0 + \Delta_\omega^0 + 1 + 3/2 = 0$. Finally, comparing the second and the fifth terms, we obtain $\Delta_w^0 + \Delta_\omega^0 + \tilde{\Delta}_\phi^0 + 2 = 0$ Now it is straightforward to find all bare scaling dimensions: $\Delta_{k_x}^0 = 3/2$, $\Delta_\omega^0 = 2$, $\Delta_\phi^0 = 1/4$, $\Delta_{\hat{\phi}}^0 = 9/4$, $\Delta_w^0 = 1/4$. (Note that the notation $\tilde{\Delta}_{\phi,\hat{\phi}}^0$ stands for the scaling dimensions of $\phi(\mathbf{k}, \omega)$ and $\hat{\phi}(\mathbf{k}, \omega)$, while $\Delta_{\phi,\hat{\phi}}^0$ denotes the scaling dimensions of $\phi(\mathbf{x}, t)$ and $\hat{\phi}(\mathbf{x}, t)$.)

According to the standard renormalization group procedure, we introduce renormalization constants: $\phi = Z_\phi^{1/2} \phi_R$, $\hat{\phi} = Z_{\hat{\phi}}^{1/2} \hat{\phi}_R$, $a = Z_a a_R$, $1/\Gamma_0 = Z_{1/\Gamma} 1/\Gamma_R$ and $\tilde{w} = Z_w \tilde{w}_R$, where

$$\tilde{w} = l^{1/4} w. \quad (57)$$

First we note that the fields ϕ and $\hat{\phi}$ have the same renormalization constants, which immediately follows from the fluctuation-dissipation theorem. Indeed, according to this theorem

$$D(x, y) = \langle \phi(x) \phi(y) \rangle \sim \text{Im} G(x, y) = \text{Im} \langle \hat{\phi}(x) \phi(y) \rangle.$$

Let us divide the Janssen-De Dominics functional into two parts:

$$J = J_R + \Delta J.$$

The first one is the "renormalized" functional $\mathcal{J}_R$

$$\begin{aligned} \mathcal{J}_R[\phi_R, \hat{\phi}_R] = & \sum_{\omega, \mathbf{k}} \hat{\phi}_{R-\omega, -\mathbf{k}} \left(a_R \frac{T}{\Gamma_R} \hat{\phi}_{R\omega, \mathbf{k}} - \left(a_R k_y^2 + \frac{k_x^2}{a_R |k_y|} \right) \phi_{R\omega, \mathbf{k}} \right. \\ & - a_R \frac{i\omega}{\Gamma_R} \phi_{R\omega, \mathbf{k}} - \tilde{w}_R l^{1/4} \frac{k_x k_y}{|k_y|} \left[\frac{\phi_R^2}{2} \right]_{\omega, \mathbf{k}} - \tilde{w}_R l^{1/4} \sum_{\Omega, \mathbf{p}} \phi_{R\omega-\Omega, \mathbf{k}-\mathbf{p}} \frac{p_x p_y}{|p_y|} \phi_{R\Omega, \mathbf{p}} \\ & \left. - \tilde{w}_R^2 a_R l^{1/2} \sum_{\Omega, \mathbf{p}} \phi_{R\omega-\Omega, \mathbf{k}-\mathbf{p}} |p_y| \left[\frac{\phi_R^2}{2} \right]_{\Omega, \mathbf{p}} \right). \end{aligned} \quad (58)$$

The second one $\Delta\mathcal{J}$ contains the counter-terms:

$$\begin{aligned}
\Delta\mathcal{J} = & \sum_{\omega,\mathbf{k}} \hat{\phi}_{R-\omega,-\mathbf{k}} \left(a_R \frac{T}{\Gamma_R} (Z_\phi Z_a Z_{1/\Gamma} - 1) \hat{\phi}_{R\omega,\mathbf{k}} \right. \\
& - \left(a_R (Z_\phi Z_a - 1) k_y^2 + (Z_\phi Z_a^{-1} - 1) \frac{k_x^2}{a_R |k_y|} \right) \phi_{R\omega,\mathbf{k}} \\
& - a_R \frac{i\omega}{\Gamma_R} (Z_\phi Z_a Z_{1/\Gamma} - 1) \phi_{R\omega,\mathbf{k}} - (Z_w Z_\phi^{3/2} - 1) \tilde{w}_R l^{1/4} \frac{k_x k_y}{|k_y|} \left[\frac{\phi_R^2}{2} \right]_{\omega,\mathbf{k}} \\
& - (Z_w Z_\phi^{3/2} - 1) \tilde{w}_R l^{1/4} \sum_{\Omega,\mathbf{p}} \phi_{R\omega-\Omega,\mathbf{k}-\mathbf{p}} \frac{p_x p_y}{|p_y|} \phi_{R\Omega,\mathbf{p}} \\
& \left. - (Z_w^2 Z_a Z_\phi^2 - 1) \tilde{w}_R^2 a_R l^{1/2} \sum_{\Omega,\mathbf{p}} \phi_{R\omega-\Omega,\mathbf{k}-\mathbf{p}} |p_y| \left[\frac{\phi_R^2}{2} \right]_{\Omega,\mathbf{p}} \right). \tag{59}
\end{aligned}$$

A simple power counting shows that all corrections are only logarithmically divergent in the dimension of $5/2$ (see section II). Hence, in what follows the ε -regularization scheme with $\varepsilon = 1/2$ is assumed.

Evaluating the diagram shown in Fig.1a up to the second order in k_y (using the bare G_0 and D_0 functions) one finds the one-loop correction to the second term of the JDF:

$$-\frac{18}{128\pi} a_R \tilde{w}_R^2 l^{1/2} \int \frac{dk_y}{k_y^{3/2}} - (Z_\phi Z_a - 1) a_R.$$

In order to cancel the divergency, we put:

$$Z_\phi Z_a = 1 - \frac{18}{128\pi} \tilde{w}_R^2 l^{1/2} \int \frac{dk_y}{k_y^{3/2}}. \tag{60}$$

The same procedure for the fourth term gives:

$$Z_a Z_\phi Z_{1/\Gamma} = 1 - \frac{1}{32\pi} \tilde{w}_R^2 l^{1/2} \int \frac{dk_y}{k_y^{3/2}}. \tag{61}$$

In the Appendix we show that both the three-leg and four-leg vertices do not have one-loop corrections. Hence:

$$Z_w Z_\phi^{3/2} = 1. \tag{62}$$

One can easily see that the one-loop correction to the term $\hat{\phi} \frac{k_x^2}{|k_y|} \phi$ in eqn (62) vanishes as well (see^{12,20}). It means that

$$Z_a^{-1} Z_\phi = 1. \quad (63)$$

Equations (60,61,62,63) have a following solution:

$$Z_a = Z_\phi = 1 - \frac{9}{128\pi} \tilde{w}_R^2 l^{1/2} \int \frac{dk_y}{k_y^{3/2}}, \quad (64)$$

$$Z_{1/\Gamma} = 1 + \frac{7}{64\pi} \tilde{w}_R^2 l^{1/2} \int \frac{dk_y}{k_y^{3/2}}, \quad (65)$$

$$Z_w = \left(1 + \frac{27}{256\pi} \tilde{w}_R^2 l^{1/2} \int \frac{dk_y}{k_y^{3/2}} \right). \quad (66)$$

Next we introduce the Gell-Mann-Low β -function $\beta = \mu \left. \frac{\partial w_R}{\partial \mu} \right|_{w, \Lambda}$ and the Callan-Symanzik anomalous dimensions $\gamma_a = \frac{\mu}{a_R} \frac{\partial a_R}{\partial \mu}$, $\gamma_\Gamma = \mu \Gamma_R \frac{\partial 1/\Gamma_R}{\partial \mu}$ and $\eta_\phi = \frac{\mu}{Z_\phi} \frac{\partial Z_\phi}{\partial \mu}$. We denote by μ the scale at which the coupling constant is equal to $\tilde{w}_R$ and denote by Λ the scale at which the coupling constant is equal to $\tilde{w}$. From (66) and the definition of Z_w , one has:

$$w_R = w \left(\frac{\Lambda}{\mu} \right)^{1/4} \left(1 - \frac{27}{128\pi} \tilde{w}^2 \Lambda^{1/2} \int_\mu^\Lambda \frac{dk_y}{k_y^{3/2}} \right). \quad (67)$$

And finally, we find the β -function, which coincide with that found in the statics case¹²:

$$\beta(\tilde{w}_R) = -\frac{1}{4} \tilde{w}_R + \frac{27}{128\pi} \tilde{w}_R^3 \quad (68)$$

The fixed point of the renormalization group flow is

$$\tilde{w}_R^{*2} = \frac{32\pi}{27}$$

After performing this procedure for anomalous dimensions one gets:

$$\gamma_a = -\frac{9}{64\pi} \tilde{w}_R^2, \quad (69)$$

$$\gamma_{1/\Gamma} = \frac{7}{32\pi} \tilde{w}_R^2, \quad (70)$$

$$\eta_\phi = \frac{9}{64\pi} \tilde{w}_R^2. \quad (71)$$

Using the result for $\tilde{w}_R^*$, it is straightforward to find $\gamma_a^* = -1/6$, $\gamma_{1/\Gamma}^* = 7/27$ and $\eta_\phi^* = 1/6$.

The long range limit of the functions G and D is found from the Callan-Symanzik equation²⁰:

$$G(t, \mathbf{x}) = \frac{1}{(x^2 + |y|^{8/3})^{7/8}} f\left(\frac{x}{|y|^{4/3}}, \frac{t}{|y|^{47/27}}\right) \quad (72)$$

$$D(t, \mathbf{x}) = \frac{1}{(x^2 + |y|^{8/3})^{1/4}} \tilde{f}\left(\frac{x}{|y|^{4/3}}, \frac{t}{|y|^{47/27}}\right) \quad (73)$$

where $f(x, y)$ and $\tilde{f}(x, y)$ are arbitrary functions.

In the static limit ($t = 0$) the exponents in the correlation function (72) are exact, as was previously found by one of the authors¹²:

$$D(\mathbf{r}) = \langle \phi(\mathbf{r}) \phi(0) \rangle \sim (x^{-1/2}, y^{-2/3}) \quad (74)$$

For the Fourier-components of the Green function eq.72, we find in the region of anomalous diffusion:

$$G(\omega, \mathbf{k}) = f_1\left(\frac{k_x}{k_y^{4/3}}, \frac{\omega}{k_y^{47/27}}\right). \quad (75)$$

The anomalous dispersion of the diffusion mode announced in the Abstract follows from the last equation. We see that the static dipole contribution is not renormalized in dynamics, as was suggested in (27,28). We also note that the exchange coupling acquires an anomalous dimension $\Delta_J = 1/3$, whereas the dynamic term ω/Γ acquires anomalous a dimension $+2/27$. Taking into account the anomalous dimension $\gamma_{1/\Gamma} = 7/27$, we conclude that the anomalous dimension of ω is $\gamma_\omega = 1/6$. The interaction between the diffusons in the scaling limit reduces the dissipation or, in other words, hardens the diffuson.

Now we can estimate the wavevector p_a , an upper boundary for anomalous diffusion. We assume that temperature is small (41). Initially, according to Eq.56, the bare vertex $w_0 = \sqrt{T}g^{1/4}/J^{3/4}$, and is also small. Under the renormalization flow, the vertex w_R grows with the inverse wavevector as the power $1/4$ (57). The RG flow starts at p_{DM} and approaches the fixed point at the root of the Gell-mann-Low function $\beta(w_R) = 0$. Invoking eq.68, we find the fixed point solution $w_R \sim 1$. Thus, the wavevecor p_a is defined as the wavevector at which $w_R \sim 1$:

$$w_0(p_{DM}/p_a)^{1/4} \sim 1. \quad (76)$$

We see that

$$p_a \sim T^2 T^{3/4}, \quad (77)$$

that is very small. Even if t is not small, $p_a \sim p_{DM} w_0^4 \sim p_0(ga/J)$. It is much smaller than p_0 .

V. SUSCEPTIBILITIES.

In this section we find the susceptibility to the magnetic field directed along the average magnetization $\langle \mathbf{S} \rangle$ (y -axis), the so-called longitudinal susceptibility. We consider the magnetic field in the form $H = H_0 + \delta H(\mathbf{x}, t)$ where H_0 is independent of $\mathbf{x}$ and t . When an additional magnetic field δH is imposed, a new vertex $\delta h_{\omega, \mathbf{k}}[\phi \hat{\phi}]_{-\omega, -\mathbf{k}}$ emerges in the JDF (26) (we denote $h = g_G \mu_B S H$; $\delta h = g_G \mu_B S \delta H$). It leads to a correction $\delta D(\omega, \mathbf{k})$ to the correlation function $D(\omega, \mathbf{k})$

$$\delta D(\mathbf{x}_1 = \mathbf{x}_2, t_1 = t_2) = \int_{\Omega} \int_{\mathbf{k}} \int_{\omega} \int_{\mathbf{q}} h(\omega, \mathbf{q}) D_0(\Omega, \mathbf{k}) [G_0(\Omega + \omega, \mathbf{k} + \mathbf{q}) + G_0^*(\Omega - \omega, \mathbf{k} - \mathbf{q})],$$

where $D_0(\omega, \mathbf{k})$ and $G_0(\omega, \mathbf{k})$ are taken from (30,31).

By definition, the susceptibility χ is:

$$\begin{aligned} \chi(\omega, \mathbf{k}, h) &= \frac{\delta}{\delta H(\omega, \mathbf{k})} \langle S_y \rangle = -\frac{1}{2} \frac{\delta}{\delta H(\omega, \mathbf{k})} \langle \phi(\mathbf{x}, t) \phi(\mathbf{x}, t) \rangle \\ &= -\frac{g_G \mu_B S}{2} \frac{\delta}{\delta h(\omega, \mathbf{k})} D(\mathbf{x}_1 = \mathbf{x}_2, t_1 = t_2). \end{aligned} \quad (78)$$

Hence,

$$\chi(\omega, \mathbf{q}) = -g_G \mu_B S \int_{\Omega} \int_{\mathbf{k}} D_0(\Omega, \mathbf{k}) G_0(\Omega + \omega, \mathbf{k} + \mathbf{q}). \quad (79)$$

In the most interesting case, when $\mathbf{q} = 0$, all integrals can be evaluated and the final answer is:

$$\begin{aligned} \chi(\omega, h) &= g_G \mu_B S \frac{\Gamma^2(3/4)}{4\pi\sqrt{\pi}} T \left(\frac{J^3}{g^2} \right)^{1/4} \frac{2\Gamma}{\omega} \\ &\quad \times \left[h^{1/4} - \left(h - \frac{i\omega}{2\Gamma} \right)^{1/4} \right]. \end{aligned} \quad (80)$$

In the limiting case $\omega = 0$, the susceptibility reads:

$$\chi = \text{const} \cdot h^{-3/4}.$$

This result has been found earlier¹⁸.

VI. CONCLUSION.

In conclusion we discuss how new modes can be observed in experiment. The new modes appear on a macroscopic scale of length of the order of magnitude $1\mu m$. Even rather weak in-plane anisotropy can suppress or disguise the new modes. Therefore, we can expect that the new dynamics will be observed in films with very weak in-plane anisotropy. The best known candidates for this role are films grown on hexagonal substrates. The hexagonal anisotropy is naturally weaker than the tetragonal one, because they are proportional to a higher degree of the relativistic parameter. Besides, the hexagonal anisotropy totally vanishes at large distances in a range of temperature from $(4/9)T_{BKT}$ till T_{BKT} , where T_{BKT} is the temperature of Berezinskii-Kosterlitz-Thouless transition^{22,29}. The simplest idea is to use the (111) face of FCC crystals, such as Ag, Au, Cu. An iron film on the (111) face of Ar has been grown by S.Bader and coworkers³⁰. Recently the Ru film has been grown on the hexagonal graphite substrate³¹. Thus, 2D ferromagnets with exact XY symmetry are available.

The next question is: what dynamic effects can be observed and at what conditions? As we have noted earlier, if the in-plane anisotropy is not especially small (less than 1K in energy scale), the only opportunity is to use a six-fold substrate in the range of temperature, not small in comparison with T_{BKT} . In this situation the observation of propagating soft modes described in section III seems to be improbable, since all they require $t \ll 1$, i.e. $T \ll T_c$. However, the anomalous diffusion can be observed even at $T \sim T_c$, given a sufficiently large scale of length ($\geq 10\mu m$). The best way to observe it is to apply a short and inhomogeneous pulse of magnetic field, and follow when the secondary signal will arrive to

fixed indicators. Such a picosecond-pulse technique has been recently used for investigation of the film dynamics³². We propose to use the same pulse technique to different films.

A quick estimation of the time $t_{x,y}$ needed for the secondary signal to reach the indicator at distances L_x and L_y along $\hat{x}$ and $\hat{y}$ directions shows: $t_{x,y} \sim (\hbar/Jp_0a)(L_{x,y}/a)^{\Delta_{x,y}^a}$, where a is the lattice constant and $\Delta_{x,y}^a$ are the anomalous-diffusion dimensions for axes $\hat{x}$ and $\hat{y}$ respectively. Thus, $t_x \sim \hbar/ga(L_x/a)^{47/36} \sim 10^{3.5}\text{sec}$ and $t_y \sim \hbar/ga(L_y/a)^{47/27} \sim 1\text{sec.}$, where we have assumed $L_x \sim L_y \sim 1\text{cm}$.

The retardation time for the secondary signal is much longer than the time for the primary signal propagation. The strong size and direction dependence of the propagation time can be used for detecting of the anomalous diffusion.

VII. ACKNOWLEDGEMENTS

The work of one of the authors (A.K.) was supported in part by the Swiss National Fond under 'Oststaaten-Soforthilfemassnahmen' grant N7GUPJ038620. Our thanks are due to M.V. Volpert for her help in preparation of the manuscript.

APPENDIX A: ONE-LOOP VERTICES CORRECTIONS

First we note that the JDF (55) has the following nontrivial symmetry¹².

$$\phi(\mathbf{k}, \omega) \rightarrow \phi(\mathbf{k}, \omega) + \varepsilon \delta^2(\mathbf{k}) \delta(\omega), \quad (\text{A1})$$

$$\hat{\phi}(\mathbf{k}, \omega) \rightarrow \hat{\phi}(\mathbf{k}, \omega), \quad (\text{A2})$$

$$k_x \rightarrow k_x - \varepsilon a w k_y, \quad (\text{A3})$$

$$k_y \rightarrow k_y. \quad (\text{A4})$$

The partition function (25) must have the same symmetry $\delta Z[j, \hat{j}] = 0$. By a standard procedure²⁰, we find the implications of the symmetry (A1-A4), known as Ward-Takahashi identities to the so-called "generating functional for proper vertices" $\Gamma[\varphi, \hat{\varphi}]$ (It is the Legendre transform of $\ln Z[j, \hat{j}]$ with respect to the fields φ and $\hat{\varphi}$):

$$\int_{\mathbf{k}} \int_{\omega} \left(\delta(\omega) \delta(\mathbf{k}) \frac{\delta \Gamma}{\delta \varphi(-\mathbf{k}, -\omega)} - aw \left[k_y \varphi(-\mathbf{k}, -\omega) \frac{\partial}{\partial k_x} \frac{\delta \Gamma}{\delta \varphi(-\mathbf{k}, -\omega)} + k_y \hat{\varphi}(-\mathbf{k}, -\omega) \frac{\partial}{\partial k_x} \frac{\delta \Gamma}{\delta \hat{\varphi}(-\mathbf{k}, -\omega)} \right] \right) = 0. \quad (\text{A5})$$

Now, writing down Taylor-like expansion of $\Gamma[\varphi, \hat{\varphi}]$ over fields φ and $\hat{\varphi}$ and plugging it into (A5), we find:

$$\Gamma_{\varphi\varphi\hat{\varphi}}(0, \bar{k}, -\bar{k}) - awk_y \frac{\partial}{\partial k_x} \Gamma_{\varphi\hat{\varphi}}(\bar{k}, -\bar{k}) = 0, \quad (\text{A6})$$

where $\bar{k}$ is used for $(\mathbf{k}, \omega)$. It was proven²⁰ that there is no corrections to the term $k_x^2/|k_y|$ due to analyticity. According to (A6), it means that corrections to the three-leg vertex vanish if we put one of the two frequencies, which the vertex depends upon, equal to zero. Hence, the corrections must depend on the product of the two ω s and are small in the framework of Section III. Considering loop-wise expansion of Γ in the same spirit, one finds:

$$\Gamma = \sum_{n=0}^{\infty} w_n \Gamma^{(n)} \approx \Gamma^{(0)} + w \Gamma^{(1)} + \dots$$

Looking only at the divergent parts of the corrections $\Gamma^{(0)}$ and $\Gamma^{(1)}$, we see that, because $\Gamma^{(0)}$ is just the bare action and has no divergencies at all, the one-loop corrections to the ω -independent three-leg vertex do not diverge. As a consequence of this fact we obtain equation (62).

APPENDIX B: CALCULATION OF SELF-ENERGY

Let us start with the Dyson Eq.(42) corresponding to a one-loop diagram on Fig.1a. We may integrate the Eq.(42) over the contour in complex plane ω such that the poles of G -function are outside the contour:

$$\begin{aligned} \Sigma_0 = \frac{p_0^2 c^6 t}{2\pi} \int f^2(\mathbf{p} - \mathbf{q}/2, \mathbf{p} + \mathbf{q}/2) & \left[\left((\Omega + \epsilon(\mathbf{p} - \mathbf{q}/2))^2 - \epsilon^2(\mathbf{p} + \mathbf{q}/2) + ib_1 \right)^{-1} \right. \\ & \left. + \left((\Omega - \epsilon(\mathbf{p} - \mathbf{q}/2))^2 - \epsilon^2(\mathbf{p} + \mathbf{q}/2) - ib_1 \right)^{-1} \right] \frac{d^2 p}{\epsilon^2(\mathbf{p} - \mathbf{q}/2)}. \end{aligned} \quad (\text{B1})$$

It is convinient to change in the second G -function $\mathbf{p}$ on $-\mathbf{p}$:

$$\Sigma_0 = \frac{p_0^2 c^6 t}{2\pi} \int \frac{f^2(\mathbf{p} - \mathbf{q}/2, \mathbf{p} + \mathbf{q}/2) d^2 p}{\Omega + \epsilon(\mathbf{p} - \mathbf{q}/2) - \epsilon(\mathbf{p} + \mathbf{q}/2) + ib_1} \left[\frac{\epsilon^{-2}(\mathbf{p} - \mathbf{q}/2)}{(\Omega + \epsilon(\mathbf{p} + \mathbf{q}/2) + \epsilon(\mathbf{p} - \mathbf{q}/2))} + \frac{\epsilon^{-2}(\mathbf{p} + \mathbf{q}/2)}{(\Omega - \epsilon(\mathbf{p} + \mathbf{q}/2) - \epsilon(\mathbf{p} - \mathbf{q}/2))} \right] \quad (\text{B2})$$

Keeping only the lowest order in small momentum $\mathbf{q}$ and frequency Ω and after a simple expansion in the brackets we find:

$$\Sigma_0 = \frac{p_0^2 c^6 t}{2\pi} \int \frac{f^2(\mathbf{p}, \mathbf{p}) d^2 p}{\Omega + \epsilon(\mathbf{p} - \mathbf{q}/2) - \epsilon(\mathbf{p} + \mathbf{q}/2) + ib_1} \times \frac{-2\Omega + 4(\epsilon(\mathbf{p} + \mathbf{q}/2) - \epsilon(\mathbf{p} - \mathbf{q}/2))}{4\epsilon^4(\mathbf{p})}. \quad (\text{B3})$$

At this point we separate the above integral into the static Ω -independent part and the rest 'dynamical' self-energy. The static Ω -independent self-energy (it always real) reads:

$$\Sigma_{st} = -\frac{p_0^2 c^6 t}{2\pi} \int \frac{f^2(\mathbf{p}, \mathbf{p})}{\epsilon^4(\mathbf{p})} d^2 p. \quad (\text{B4})$$

Now let us take into account the static self-energy given by the diagram on Fig.1b:

$$\Sigma_b = \frac{p_0 c^4 t}{4\pi} \int \frac{p_y^2 - p_x^2}{p^2 \epsilon^2(\mathbf{p})} d^2 p \quad (\text{B5})$$

Comparing Eq.(B4) (remember that $f(\mathbf{p}, \mathbf{p}) = p_x^2 p_y^2 / p^2$) and Eq.(B5) we conclude that these cancel each other in the $\mathcal{A}$ -shell. Thus, we have verified explicitly that the $\Omega, \mathbf{q}$ -independent part of the self-energy is strictly zero as guaranteed by the Ward identity due to the rotation symmetry of the system (see Appendix A). To get non-zero static self-energy we have to expand self-energies like Eqs.(B4,B5) in powers of the transferred momentum $\mathbf{q}$. This will be done in the Section IV using statics field-theoretical technic not dynamical as in this Appendix. The result is that $\mathbf{q}$ -dependent static self-energy only matters when the spin-wave momentum $\mathbf{q}$ is so small that anomalous diffusion sets up. On shorter wavelengths like $q < p_{DM}$ we may safely neglect the contribution of Eq.(B4,B5).

Now we return to the Ω -dependent dynamical part of the self energy:

$$\Sigma_0(\chi, q) = \frac{p_0^2 c^6 t}{2\pi} 2\Omega \int \frac{\sin^2(2\phi)}{\Omega - q \cos \phi + ib_1} \frac{dp d\phi}{4c^4 p} \quad (\text{B6})$$

The integral over p gives exactly the logarithmic factor L :

$$\Sigma_0(\chi, q) = \frac{p_0^2 c^2 t}{\pi} L \Omega \int \frac{\sin^2 \phi \cos^2 \phi}{\Omega - q \cos \phi + ib_1} d\phi \quad (\text{B7})$$

The imaginary part of Eq.(B7) could be easily found in the limit $\Omega, cq \gg b_1$. In this case we use the formula

$$\frac{1}{x - i0} = P \frac{1}{x} + i\pi \delta(x)$$

to find the $\Im$ -part of Eq.(46). The $\Re$ -part is also simple to calculate:

$$\Re \Sigma_0(\chi, q) = \frac{p_0^2 c^2 t}{\pi} L \frac{\Omega}{q} \int \frac{\sin^2 \phi \cos \phi [(-\Omega + q \cos \phi) + \Omega]}{\Omega - q \cos \phi} d\phi, \quad (\text{B8})$$

and then simplify:

$$\Re \Sigma_0(\chi, q) = \frac{p_0^2 c^2 t}{\pi} L \frac{\Omega^2}{q} \int \frac{\sin^2 \phi \cos \phi}{\Omega - q \cos \phi} d\phi. \quad (\text{B9})$$

We repeat the same step:

$$\Re \Sigma_0(\chi, q) = \frac{p_0^2 c^2 t}{\pi} L \frac{\Omega^2}{q^2} \int \frac{\sin^2 \phi (-\Omega + q \cos \phi + \Omega)}{\Omega - q \cos \phi} d\phi \quad (\text{B10})$$

with further simplification:

$$\Re \Sigma_0(\chi, q) = \frac{p_0^2 c^2 t}{\pi} L \frac{\Omega^3}{q^2} \int \frac{\sin^2 \phi}{\Omega - q \cos \phi} d\phi - \frac{p_0^2 c^2 t}{\pi} L \frac{\Omega^2}{q^2} \frac{2\pi}{2} \quad (\text{B11})$$

One could easily continue the same procedure to find

$$\Re \Sigma_0(\chi, q) = p_0^2 c^2 t L \frac{\Omega^2}{q^2} (2 \frac{\Omega^2}{q^2} - 1) = p_0^2 c^2 t L \cos^2 \chi \cos(2\chi), \quad (\text{B12})$$

which is exactly the Eq.(46) of the III Section.

Finally let us show that the self-energy in the $\mathcal{A}$ -shell is negligible. We first assume that it does negligible: $a^2(\omega, \mathbf{p}) \ll \epsilon^2(\mathbf{p})$. Then we consider the self-energy Eq.(B2), provided the external frequency and momentum lies on the mass-shell inside the $\mathcal{A}$ -shell: $\Omega = \epsilon(\mathbf{q})$. Let also $\sqrt{\lambda/J} \gg q < p_0$. One could easily verify that the main contribution comes if $q \ll p$. In this case we may use the Eq.(B6):

$$\Sigma_0 = \frac{p_0^2 c^6 t}{2\pi} 2cq \int \frac{\sin^2(2\phi)}{c\sqrt{q^2 + p_0 q_x^2/q - q \cos \phi + ib_1}} \frac{dp d\phi}{4c^4 p} \quad (\text{B13})$$

Integration over p gives $\log(\sqrt{\lambda/J}/q)$, whereas the integration over the relative direction of internal and external spin-waves gives factor $\sqrt{q^3/p_0 q_x^2}$. Thus, the real self-energy

$$a^2(\Omega, \mathbf{q}) \sim tLc^2p_0\sqrt{p_0q}.$$

At small t this result justifies our neglecting of self-energy in the $\mathcal{A}$ -shell.

APPENDIX C: THE ANGULAR DEPENDENCE OF THE SELF-ENERGY OPERATOR.

In this appendix we analyze the angular dependence of the self-energy part found in Section III. Repeating all arguments, one can find that eq. (45) must be written in slightly modified form:

$$\Sigma_0(\Omega, q, \phi) = \frac{c^2 p_0^2 t}{4\pi} \int \frac{c^4 p^3 dp}{\epsilon^4(\mathbf{p})} \frac{\Omega \sin^2(2\psi + 2\phi) d\psi}{\Omega - cq \cos \psi + ib_1}. \quad (\text{C1})$$

Now again, assuming $cq \gg b_1$, one finds:

$$\Sigma_0(r, \phi) = c^2 p_0^2 t L \left[r^2 (2r^2 - 1 - 2r\sqrt{r^2 - 1}) \cos(4\phi) + \frac{1}{2} \frac{r}{\sqrt{r^2 - 1}} \sin^2(2\phi) \right], \quad (\text{C2})$$

where $r = \Omega/cq$. In the case $\phi = 0$ and $r = \cos \chi$ the formula (46) is recovered.

The main contribution to the integral in eq. (49) comes from a region of very small angles ϕ . Hence, the correction to the Σ_0 we have found above is not important in this calculations and equation (50) still holds.

By the next step we need to plug $b_1(\psi)$ from (50) to the nonzero-angle form of equation (48)

$$\Sigma(\Omega, q, \phi) = i\Omega c^2 p_0^2 t L \int_0^{2\pi} \frac{d\psi}{4\pi} \frac{\sin^2(2\psi + 2\phi)}{ib_1(\psi) + \Omega - cq \cos \psi}. \quad (\text{C3})$$

Evaluating this integral, one finds:

$$\Sigma(|\Omega| \ll cq) = i\Omega c\tilde{p} \left[\cos^2(2\phi) - \alpha \left(\frac{q}{p_{DM}} \right)^{-3/5} \sin^2(2\phi) \right] \quad (\text{C4})$$

$$\Sigma(|\Omega| \gg cq) = i\Omega c\tilde{p} \left[\cos^2(2\phi) - \alpha \left(\frac{|\Omega|}{cp_{DM}} \right)^{-3/5} \sin^2(2\phi) \right], \quad (\text{C5})$$

where we denote $\tilde{p} = \beta_0 t^{1/4} L^{5/4} p_0$, $p_{DM} = \beta_1 t^{3/4} L^{-1/4} p_0$ and $\alpha = 2^{-1/5}/\beta_0\beta_1 \approx 0.54$. In what follows we will also use the notation $p_c = p_0 \sqrt{tL}$. While equation (C2) for Σ holds for $cq, \Omega \gg b_1(\psi)$, equations (C4) and (C5) are valid in the opposite case $cq, \Omega \ll b_1(\psi)$.

Now we can analyze the dispersion relation $\omega^2(\mathbf{q}) = \epsilon^2(\mathbf{q}) + \Sigma(\omega, \mathbf{q})$ more accurately. In experiment usually $t \ll 1$ so $p_0 \gg \tilde{p} \gg p_c \gg p_{DM}$. Easy, but tedious calculations show that there can exist up to 9 asymptotic regions in the momentum space with different dispersion relations.

$$\begin{aligned}
\omega^2 &= c^2 p_0 q \sin^2 \phi, & \text{if } p_{DM} \frac{\sin^{3/2}(2\phi) \sin(\phi/2)}{\cos \phi} \ll q \ll p_0 \sin^2 \phi \\
\omega^2 &= \frac{1}{2} c^2 q^3 \frac{p_0}{p_c^2} \phi^2, & \text{if } \frac{p_c^2}{p_0} \phi^2 \left(1 + \frac{p_0 p_{DM}}{p_c^2} \phi^{1/2}\right) \ll q \ll \frac{p_c^2 p_0 \phi^2}{p_0^2 \phi^4 + p_c^2} \\
\omega^2 &= \frac{1}{2} c^2 q^3 \frac{p_0}{p_c^2}, & \text{if } p_{DM} (\pi/2 - \phi)^{1/2} \ll q \ll \frac{p_c^2}{p_0} \\
\omega^2 &= \frac{c^2}{2} \frac{q^4}{p_c^2}, & \text{if } p_0 \phi^2 \ll q \ll p_c \\
\omega &= -icq^2 \frac{p_0}{p_c^2}, & \text{if } p_{DM} \phi^{5/2} \ll q \ll \frac{p_c^2}{p_0} \phi^2 \\
\omega &= -icq \frac{p_0}{\tilde{p}} \phi^2, & \text{if } p_{DM} \phi^{10/3} \ll q \ll p_{DM} \phi^{5/2}; \quad \phi \ll \sqrt{\tilde{p}/p_0} \\
\omega &= -icq \frac{p_0}{\tilde{p}} \phi^2, & \text{if } p_{DM} \phi^{4/3} \ll q \ll p_{DM} \phi^{1/2}; \quad \sqrt{\tilde{p}/p_0} \ll \phi \ll 1 \\
\omega &= -icq \frac{p_0}{\tilde{p}}, & \text{if } \frac{\tilde{p} p_{DM}}{p_0} (\pi/2 - \phi)^{10/3} \ll q \ll \frac{\tilde{p} p_{DM}}{p_0} (\pi/2 - \phi)^{1/2} \\
\omega^{2/5} &= -iq \frac{c^{2/5} p_0}{4\alpha \tilde{p} p_{DM}^{3/5}} \frac{1}{\cos^2 \phi}, & \text{if } p_{DM} (\frac{\tilde{p}}{p_0})^{5/3} (\pi/2 - \phi)^{10/3} \ll q \ll \frac{\tilde{p} p_{DM}}{p_0} \phi^{4/3} (\pi/2 - \phi)^{10/3}
\end{aligned}$$

where $0 < \phi < \pi/2$

In figure 3 we show the case $t \sim 0.3$, $L \sim 1$ and $p_0 = 1$.

REFERENCES


- ¹ A.A.Abrikosov, L.P.Gor'kov and I.E.Dzyaloshinsky, *Methods of Quantum Field Theory in Statistical Physics*, New York, 1968.
- ² A.B.Migdal, ZhETF **34**, 417 (1958)
- ³ D.V.Khveshchenko and P.C.E.Stamp, Phys.Rev.Lett. **71**, 2118 (1993); B.L.Altshuler, L.B.Ioffe, and A.Millis, Phys. Rev. B **51**, (1994);
- ⁴ A.Stern and B.I.Halperin, Phys.Rev.B, **52**, 5890 (1995)
- ⁵ K.Kawasaki, Phys. Rev. **150**, 291 (1966)
- ⁶ L.P.Kadanoff and J.Swift, Phys. Rev. **166**, 89 (1968)
- ⁷ P.C. Hohenberg and B.I. Halperin, Rev. Mod. Phys. **49**, 435 (1977)
- ⁸ R.Zeyher, Ferroelectrics, **66**, 217 (1986).
- ⁹ E.I.Kats and V.V.Lebedev, *Fluctuational effects in the dynamics of liquid crystals* (Springer-Verlag 1993).
- ¹⁰ G.Grüner, Rev.Mod.Phys., **60**,1129 (1988).
- ¹¹ E.Frey and F.Schwabl, Adv.Phys. **43**, 577 (1994)
- ¹² A.Kashuba, Phys.Rev.Lett. **73**, 2264 (1994)
- ¹³ P.C.Martin, E.D.Siggia and H.A.Rose, Phys.Rev.A, **8**, 423 (1973); C.De Dominicis and L.Peliti, Phys.Rev.B, **18**, 353 (1978); H.K.Janssen, Z.Phys.B, **23**, 372 (1976)
- ¹⁴ A.Kashuba, Ar.Abanov and V.L.Pokrovsky, cond-mat/9603160 to be published in Phys.Rev.Lett.
- ¹⁵ A.V. Chubukov and M.I. Kaganov, Soviet Physics-Uspekhi **30**, 1015 (1987)
- ¹⁶ The classical exchange constant J and anisotropy λ used in this paper differs from that

- in quantum Heisenberg model: $J_{\text{cl}} = J_{\text{qua}} S^2$ and $\lambda_{\text{cl}} = \lambda_{\text{qua}} S^2$
- ¹⁷ A.Kashuba and V.L.Pokrovsky, Phys.Rev.B **48**, 10335 (1993)
- ¹⁸ S.V.Maleev, ZhETF **71**, 2375 (1976); V.L.Pokrovsky and M.V.Feigelman, ZhETF **72**, 557 (1977)
- ¹⁹ N.D.Mermin and H.Wagner, Phys.Rev.Lett. **17**, 1133 (1966)
- ²⁰ J. Zinn-Justin, *Quantum Field Theory and Critical Phenomena*. Oxford, 1993.
- ²¹ L.D.Landau and E.M.Lifshitz, *Electrodynamics of Continuous Media*.
- ²² A.Ya. Blank, V.L. Pokrovsky and G.V. Uimin, Journal of Low Temperature Physics.
- ²³ V.L. Berezinsky, ZhETP **59**, 907 (1970) [Sov.Phys.JETP **32**, 493 (1971)]
- ²⁴ An allowed by symmetry term $\sim R' \dot{\pi} \ddot{\pi}$ in the dissipative function is of higher order in time derivatives, since $\dot{\pi} \sim \ddot{\phi}$. dissipation functional are also irrelevant.
- ²⁵ L.D.Landau and E.M.Lifshitz, *Statistical Physics*. Pergamon Press.
- ²⁶ The a -function is negligible in the $\mathcal{A}$ -shell. If one assumes that $a^2(\omega, \mathbf{p}) \ll \epsilon^2(\mathbf{p})$ inside the $\mathcal{A}$ -shell, then from equation (42) one finds: $a^2(\epsilon(\mathbf{q}), \mathbf{q}) \sim t L c^2 p_0 \sqrt{p_0 q}$, justifying the assumption.
- ²⁷ A.Aharony, in *Phase Transition and Critical Phenomena*, vol.6, eds. C.Domb and M.S.Green (Academic press 1976).
- ²⁸ R.A.Pelcovits and B.I.Halperin, Phys.Rev.B **19**, 4614 (1979).
- ²⁹ J.V.Jose, L.P. Kadanoff, S. Kirkpatrick, and D.R. Nelson, Phys.Rev.B **16**, 1217 (1977).
- ³⁰ Z.Q. Qiu, J. Pearson, and S.D. Bader, Phys. Rev. Lett. **67**, 1646 (1991).
- ³¹ R. Pfandzelter, G. Steierl, and C.Rau, Phys. Rev. Lett. **74**, 3467 (1995).
- ³² H.C. Siegmann, E.L.Garwin, C.Y. Prescott, J. Feldman, D. Mauri, D. Weller, R. Al-


lenspach, and W. Weber, Journ. Magn. Magn. Mat. **151**, L8 (1955).

Figure Caption


- FIG.1. a,b) The main contribution to the self-energy. The functions G and D are given by (34). The three-leg vertices in fig.1.a and the four-leg vertex in fig.1.b are from (32). c) Two-loop correction to the self-energy. Momenta of internal lines are indicated.
- FIG.2. The most 'dangerous' two-loop diagram.
- FIG.3. Maps of the regions with different dispersion relations.



$G(\omega, p)$



$D(\omega + \Omega, p + q)$



D

